


Introduction to Hierarchical Linear Modeling with R




An Introduction to HLM with R

Dr. J. Kyle Roberts

First Things First

- Robinson (1950) and the problem of contextual effects
- The “Frog-Pond” Theory



An Introduction to HLM with R

Dr. J. Kyle Roberts

A Brief History of Multilevel Models

- Nested ANOVA designs
- Problems with the ANCOVA design
 - “Do schools differ” vs. “Why schools differ?”
 - ANCOVA does not correct for intra-class correlation (ICC)

An Introduction to HLM with R

Dr. J. Kyle Roberts

3

Strengths of Multilevel Models

- Statistical models that are not hierarchical sometimes ignore structure and report underestimated standard errors
- Multilevel techniques are more efficient than other techniques
- Multilevel techniques assume a general linear model and can perform all types of analyses

An Introduction to HLM with R

Dr. J. Kyle Roberts

4

Multilevel Examples

- Students nested within classrooms
- Students nested within schools
- Students nested within classrooms within schools
- Measurement occasions nested within subjects (repeated measures)
- Students cross-classified by school and neighborhood
- Students having multiple membership in schools (longitudinal data)
- Patients within a medical center
- People within households

An Introduction to HLM with R

Dr. J. Kyle Roberts

5

Do we really need HLM/MLM?

- “All data are multilevel!”
- The problem of independence of observations
- The “inefficiency” of OLS techniques


An Introduction to HLM with R

Dr. J. Kyle Roberts

6

Differences in HLM and Other Methods

- HLM is based on Maximum Likelihood and Empirical Bayesian estimation techniques
- $1 + 1 = 1.5$




An Introduction to HLM with R

Dr. J. Kyle Roberts

7

Graphical Example of Multilevel ANOVA



An Introduction to HLM with R

Dr. J. Kyle Roberts

8

Notating the HLM ANOVA

- The full model would be:

$$y_{ij} = \gamma_{00} + u_{0j} + e_{ij}$$
 - Level-1 model is:

$$y_{ij} = \beta_{0j} + e_{ij}$$
 - Level-2 model is:


$$\beta_{0j} = \gamma_{00} + u_{0j}$$
- $$\left\{ \begin{array}{l} y_{11} = \beta_1 + e_{11} \\ y_{21} = \beta_1 + e_{21} \\ \dots \\ y_{ij} = \beta_j + e_{ij} \end{array} \right.$$
- $$\left\{ \begin{array}{l} \beta_1 = \gamma_{00} + u_1 \\ \beta_2 = \gamma_{00} + u_2 \\ \dots \\ \beta_j = \gamma_{00} + u_j \end{array} \right.$$

An Introduction to HLM with R

Dr. J. Kyle Roberts

9

Understanding Errors




An Introduction to HLM with R

Dr. J. Kyle Roberts

10

Fixed vs. Random Coefficients

- Fixed Slopes and Intercepts




An Introduction to HLM with R

Dr. J. Kyle Roberts

11

Fixed vs. Random Coefficients

- Fixed Slopes and Intercepts
- Random Intercepts and Fixed Slopes




An Introduction to HLM with R

Dr. J. Kyle Roberts

12

Fixed vs. Random Coefficients

- Fixed Slopes and Intercepts
- Random Intercepts and Fixed Slopes
- **Fixed Intercepts and Random Slopes**




An Introduction to HLM with R

Dr. J. Kyle Roberts

13

Fixed vs. Random Coefficients

- Fixed Slopes and Intercepts
- Random Intercepts and Fixed Slopes
- Fixed Intercepts and Random Slopes
- **Random Slopes and Intercepts**



An Introduction to HLM with R

Dr. J. Kyle Roberts

14

Let's Give This A Shot!!!

- An example where we use a child's level of "urbanicity" (a SES composite) to predict their science achievement
- Start with Multilevel ANOVA (also called the "null model")

$$science_{ij} = \gamma_{00} + u_{0j} + r_{ij}$$

Grand mean Group deviation Individual diff.

An Introduction to HLM with R

Dr. J. Kyle Roberts

15

Intraclass Correlation

- The proportion of total variance that is *between* the groups of the regression equation
- "The degree to which individuals share common experiences due to closeness in space and/or time" Kreft & de Leeuw, 1998.
- a.k.a – ICC is the proportion of group-level variance to the total variance
- LARGE ICC DOES NOT EQUAL LARGE DIFFERENCES BETWEEN MLM AND OLS (Roberts, 2002)
- Formula for ICC:
$$\rho = \frac{\sigma_{u0}^2}{\sigma_{u0}^2 + \sigma_{e0}^2}$$

An Introduction to HLM with R

Dr. J. Kyle Roberts

16

Statistical Significance???

- Chi-square vs. degrees of freedom in determining model fit
- The problem with the df
- Can also compute statistical significance of variance components (only available in some packages)

An Introduction to HLM with R

Dr. J. Kyle Roberts

17

The Multilevel Model – Adding a Level-1 Predictor

- Consider the following 1-level regression equation:
 - $y = a + bx + e$
 - y = response variable
 - a = intercept
 - b = coefficient of the response variable (slope)
 - x = response variable
 - e = residual or error due to measurement

An Introduction to HLM with R

Dr. J. Kyle Roberts

18


The Multilevel Model (2)

- The fixed coefficients multilevel model is a slight variation on the OLS regression equation:
 - $y_{ij} = a + bx_{ij} + u_j + e_{ij}$
 - Where “i” defines level-1, “j” defines level-2, u_j is the level-2 residual and e_{ij} is the level-1 residual
- Using slightly different annotation we can transform the above equation to:
 - $y_{ij} = \gamma_{00} + \gamma_{10}x_{ij} + u_{0j} + e_{ij}$
 - Where γ_{00} now defines the constant/intercept “a” and γ_{10} defines the slope

The Multilevel Model (3)

- From the previous model:
 - $y_{ij} = \gamma_{00} + \gamma_{10}x_{ij} + u_{0j} + e_{ij}$
- We can then transform this model to:
 - $y_{ij} = \beta_{0j} + \beta_{1j}x_{1ij} + e_{ij}$ ← Level-1 Model
 - $\beta_{0j} = \gamma_{00} + u_{0j}$ ← Level-2 Model
 - $\beta_{1j} = \gamma_{10}$ ←
 - With variances $u_{0j} = \sigma_{u0}^2$ $e_{ij} = \sigma_{eij}^2$

Understanding Errors



An Introduction to HLM with R

Dr. J. Kyle Roberts

21

Adding a Random Slope Component

- Suppose that we have good reason to assume that it is inappropriate to “force” the same slope for “urbanicity” on each school
 - Level-1 Model $\rightarrow y_{ij} = \beta_{0j}x_0 + \beta_{1j}x_{1ij} + r_{ij}$
 - Level-2 Model $\rightarrow \beta_{0j} = \gamma_{00} + u_{0j}$
 $\beta_{1j} = \gamma_{10} + u_{1j}$


Complete Model $\rightarrow science_{ij} = \gamma_{00} + u_{0j} + (\gamma_{10} + u_{1j})urban + r_{ij}$

An Introduction to HLM with R

Dr. J. Kyle Roberts

22

Understanding Errors Again



An Introduction to HLM with R

Dr. J. Kyle Roberts

23

Model Fit Indices

- Chi-square $-2 * \ell$
- Akaike Information Criteria

$$AIC = -2 * \ell + 2K$$

- Bayesian Information Criteria

$$BIC = -2 * \ell + K * \ln(N)$$

An Introduction to HLM with R

Dr. J. Kyle Roberts

24

To Center or Not to Center

- In regression, the intercept is interpreted as the expected value of the outcome variable, when all explanatory variables have the value zero
- However, zero may not even be an option in our data (e.g., Gender)
- This will be especially important when looking at cross-level interactions
- General rule of thumb: If you are estimating cross-level interactions, you should grand mean center the explanatory variables.

An Introduction to HLM with R

Dr. J. Kyle Roberts

25

An Introduction to R

An Introduction to HLM with R

Dr. J. Kyle Roberts

26

R as a Big Calculator

- Language was originally developed by AT&T Bell Labs in the 1980's
- Eventually acquired by MathSoft who incorporated more of the functionality of large math processors
- The commands window is like a big calculator

```
> 2+2           > 3*5+4
[1] 4           [1] 19
```

Object Oriented Language

- A Big Plus for R is that it utilizes object oriented language.

```
> x<-2+4
> x
[1] 6

> y<-3+5
> y
[1] 8

> x+y
[1] 14

> x<-1:10
> x
[1] 1 2 3 4 5 6
7 8 9 10
> mean(x)
[1] 5.5

> 2*x
[1] 2 4 6 8 10 12 14 16
18 20
> x^2
[1] 1 4 9 16 25 36
49 64 81 100
```

Utilizing Functions in R

- R has many “built in” functions (c.f., “Language Reference” in the “Help” menu)
- Functions are commands that contain “arguments”
- seq function has 4 arguments
 - seq(from, to, by, length.out, along.with)

```
> ?seq
> seq(from=1, to=100, by=10)
[1] 1 11 21 31 41 51 61 71
81 91
> seq(1, 100, 10)
> seq(1, by=10,
length=4)
[1] 1 11 21 31
```

An Introduction to HLM with R

Dr. J. Kyle Roberts

29

Making Functions in R

```
> squared<-function(x){x^2}
> squared(5)
[1] 25

> inverse<-function(x){1/x}
> num<-c(1,2,3,4,5)
> inverse(num)
[1] 1.0000000 0.5000000 0.3333333
0.2500000 0.2000000
```

An Introduction to HLM with R

Dr. J. Kyle Roberts

30

Reading a Dataset

```
> example<-read.table(file, header=T)
> example<-read.table("c:/aera/example.txt", header=T)
> head(example)
```

Other Software Packages for HLM Analysis

- Good Reviews at <http://www.cmm.bristol.ac.uk/>
 - MLwiN
 - SAS – PROC MIXED, PROC NL MIXED, GLIMMIX
 - S-PLUS – lme, nlme, glme
 - *Mplus*
 - SPSS – Version 12 and later
 - STATA